

From carbon nanostructures to 2D semiconductors beyond graphene*

David Tománek

Physics and Astronomy Department, Michigan State University, East Lansing, MI 48824, USA

tomanek@pa.msu.edu • <http://www.pa.msu.edu/people/tomanek>

Abstract

If graphene had a band gap, it would probably be the optimum 2D system for electronics applications. While graphene research is still revealing intriguing properties of this unusual material, a new class of 2D semiconductors has been rapidly gaining attention, namely layered black phosphorus and related phosphorene monolayers [1]. These 2D systems display a tunable, direct fundamental band gap and thus are ideal candidates for optoelectronics applications. Recent Quantum Monte Carlo (QMC) calculations show that the inter-layer bonding, while weak, is not well described by dispersive van der Waals (vdW) interactions [2]. As seen in Fig. 1, QMC results differ qualitatively from vdW-enhanced DFT functionals and the common designation of similar systems as “van der Waals solids” is strictly incorrect. Also other group V systems including monolayers of $\text{As}_x\text{P}_{1-x}$ [3], IV-VI compounds such as SiS [4] with the same average valence, and related 2D phosphorus carbide [5] share the same nonplanarity of their structure with phosphorene. These systems share another similarity with phosphorene, namely the dependence of the fundamental band gap on the number of layers and in-layer strain. A surprising twist on the story of group V semiconductors is the recent discovery of yet another phosphorus allotrope, namely a quasi-1D coil structure [6,7] that is the most stable phosphorus allotrope known to date. Predictive *ab initio* calculations provide here a useful guidance to experimental studies.




FIG.1. Calculated electron density difference $\Delta\rho = \rho_{\text{tot}}(\text{bulk}) - \sum \rho_{\text{tot}}(\text{monolayers})$ representing the charge redistribution caused by assembling the bulk structure from isolated monolayers. (a) Diffusion Monte Carlo (DMC) isosurfaces bounding regions of excess electron density (dark brown) and electron deficiency (light brown), with respective values $\pm 6.5 \times 10^{-3} \text{ e}/\text{\AA}^3$. (b) $\langle \Delta\rho(z) \rangle$ for DMC and selected DFT functionals averaged across the x - y plane of the layers, with z/c indicating the relative position of the plane in the unit cell. (From Ref. [2]).

* Partly supported by the NSF/AFOSR EFRI 2-DARE grant number #EFMA-1433459.

References:

- [1] H. Liu *et al.* [ACS Nano 8, 4033 \(2014\)](#).
- [2] L. Shulenburger, A.D. Baczewski, Z. Zhu, J. Guan, and D. Tománek, [Nano Lett. 15, 8170 \(2015\)](#).
- [3] Zhen Zhu, Jie Guan, and David Tománek, [Nano Lett. 15, 6042 \(2015\)](#).
- [4] Zhen Zhu, Jie Guan, Dan Liu, and David Tománek, [ACS Nano 9, 8284 \(2015\)](#).
- [5] Jie Guan, Dan Liu, Zhen Zhu, and David Tománek, [Nano Lett. 16, 3247 \(2016\)](#).
- [6] Dan Liu, Jie Guan, Jingwei Jiang, and David Tománek, [Nano Lett. 16, 7865 \(2016\)](#).
- [7] Jinying Zhang *et al.* [Angew. Chem. Int. Ed. 56, 1850 \(2017\)](#).